

MATHEMATICS

ON THE SIMULTANEOUS REPRESENTATION OF A GIVEN PAIR OF INTEGERS AS THE SUM RESPECTIVELY OF FOUR INTEGERS AND THEIR SQUARES. I.

BY

LILY SESHU

(Communicated by Prof. H. D. KLOOSTERMAN at the meeting of September 24, 1960)

Acknowledgement

The author wishes to thank Professor PAUL T. BATEMAN for suggesting the problem and for the many profitable discussions she had with him.

Introduction

In this paper an expression is obtained for the arithmetic function $r_s(n, m)$ which represents the number of solutions of the diophantine equations:

$$(a) \quad x_1^2 + x_2^2 + \dots + x_s^2 = n,$$

$$(b) \quad x_1 + x_2 + \dots + x_s = m,$$

when the value of s is four.

KLOOSTERMAN used the circle method of Hardy and Littlewood to obtain heuristically a function $\varrho_s(n, m)$ which approximates $r_s(n, m)$. When $3 \leq s \leq 8$, it turns out that $\varrho_s(n, m) = r_s(n, m)$. KLOOSTERMAN himself [1] showed this for $s=3, 5, 7$ and BRONKHORST [1] did the same for $s=6, 8$. Van der Blij used an algebraic method to obtain an exact formula for $r_4(n, m)$. LOMADZE [1] on the other hand summed the singular series to obtain a closed expression for $\varrho_s(n, m)$ for $s \geq 3$. His results together with those of Van der Blij indicate that $r_4(n, m) = \varrho_4(n, m)$.

In this paper the same result is proved directly, utilizing the technique Bronkhorst and Kloosterman (loc. cit.). A convergence problem arises in the process similar to that encountered by several other authors in the solution of problems of a similar nature. See for instance KLOOSTERMAN (loc. cit.), H. STREEFKERK [1], P. T. BATEMAN [1], and some others. The technique used to overcome the difficulty, namely the introduction of a convergence-producing factor into the terms of a formal series, goes back to HECKE [1].

Some symbols used

Let a, b, q, x be integers.

$$e\left(\frac{a}{q}\right) : e^{2\pi i a/q}.$$

$$\sum_{a(q)} f(a) \text{ or } \sum_{a(\bmod q)} f(a) : \sum_{a=0}^{q-1} f(a).$$

$$\sum'_{a(q)} f(a) \text{ or } \sum'_{a \bmod q} f(a) : \sum_{\substack{a=0 \\ (a,q)=1}}^{q-1} f(a).$$

If $a \neq 0$,

$\left(\frac{a}{b}\right)$: Jacobi or Kronecker symbol as appropriate, ($b > 1$).

$$\left(\frac{a}{1}\right) = 1.$$

$$\sum_m f(m) : \sum_{m=-\infty}^{\infty} f(m).$$

$$S(a, q, b) : \sum_{x \bmod q} e\left(\frac{ax^2 + bx}{q}\right).$$

Section 1. *Definitions and basic properties.*

In this section the functions $\theta_z(v|\tau)$ and $\theta(v|\tau)$ are defined. $\theta_z(v|\tau)$ is defined by means of a double Laurent series in $e^{2\pi iv}$ and $e^{\pi i \tau}$ where v, τ and z are complex variables with $\mathcal{I}(\tau) > 0$ — the coefficient of $e^{2\pi i v m} e^{\pi i \tau n}$ being $\varrho_4(n, m)$. In the initial definition of $\theta_z(v|\tau)$, z is restricted to the half plane $\mathcal{R}(z) > 1/2$. Using a certain transformation on $\theta_z(v|\tau)$, it is shown that the latter may be continued analytically into the region $\mathcal{R}(z) > -1/2$ and that $\theta(v|\tau)$ is identical with the value of $\theta_z(v|\tau)$ at $z=0$.

Since the lines of work are necessarily similar to those of BRONKHORST [1] and KLOOSTERMAN [1], every attempt has been made to retain the same notation. For the same reason the subscript s is used rather than four, even though in the whole of this paper s is assumed to be four unless specifically otherwise stated.

We need certain results concerning $\varrho_s(n, m)$ and the singular series prior to defining $\theta(v|\tau)$.

From the work of KLOOSTERMAN [1, p. 325, p. 333] we obtain an expression for $\varrho_s(n, m)$. Let

$$\Delta = ns - m^2.$$

If $\Delta \leq 0$, we put:

$$(1) \quad r_s(n, m) = \varrho_s(n, m) = \begin{cases} 0 & \text{for } \Delta < 0 \text{ or } s \nmid m \\ 1 & \text{for } \Delta = 0 \text{ and } s|m. \end{cases}$$

If $\Delta > 0$, we put

$$(2) \quad \varrho_s(n, m) = \frac{\pi^{(s-1)/2} \Delta^{(s-3)/2}}{s^{(s-2)/2} \Gamma\left(\frac{s-1}{2}\right)} \sum_{q=1}^{\infty} \sum'_{a(q)} \sum_{b(q)} \left(\frac{S(a, q, b)}{q}\right)^s e\left(\frac{-na - mb}{q}\right).$$

In future we denote by η_s the factor

$$\frac{\pi^{(s-1)/2}}{s^{(s-2)/2} \Gamma\left(\frac{s-1}{2}\right)}.$$

Then $\varrho_s(n, m)$ is related to the singular series $\mathfrak{S}_s(n, m)$ thus:

$$(3) \quad \varrho_s(n, m) = \eta_s \mathfrak{S}_s(n, m) \cdot \Delta^{(s-3)/2}.$$

If we define:

$$(4) \quad B_q(n, m) = \sum'_{a(q)} \sum_{b(q)} \left(\frac{S(a, q, b)}{q} \right)^s e\left(\frac{-na - mb}{q}\right),$$

then we have:

$$\varrho_s(n, m) = \eta_s \sum_{q=1}^{\infty} B_q(n, m) \cdot \Delta^{(s-3)/2}.$$

KLOOSTERMAN shows [1, p. 323 and p. 333] that

$$(5) \quad B_{q_1 q_2}(n, m) = B_{q_1}(n, m) \cdot B_{q_2}(n, m),$$

provided $(q_1, q_2) = 1$.

G. A. LOMADZE [1] considers the convergence and effects the summation of the singular series for $s \geq 3$. $\mathfrak{S}_s(n, m)$ is absolutely convergent only for $s \geq 5$ but converges for $s = 3, 4$. Lomadze's results for $s = 4$ are stated in equations (7a) to (9) below. These equations yield the following observations: If q is odd, $B_q(n, m)$ depends only on $\Delta = ns - m^2$. If q is even but $2^4 \nmid q$, $B_q(n, m)$ depends on the residue classes modulo 4 (in some cases, modulo 8) to which n and m belong. If $2^4 | q$, $B_q(n, m)$ depends on Δ and the residue classes of m modulo 4.

In what follows the argument of B_q may be dropped if no ambiguity results.

If h is a prime, we define:

$$(6) \quad \chi_h(n, m) = 1 + B_h + B_{h^2} + \dots$$

For the rest of this paper p denotes an odd prime.

Lomadze's results for $B_q(n, m)$ when $s = 4$:

$$q = 2^\lambda, \quad 1 \leq \lambda \leq 3. \quad (i)$$

$$(7a) \quad B_2 = (-1)^{n+m}$$

$$(7b) \quad B_{2^2} = \begin{cases} 0 & \text{if } 2 \nmid m, \text{ or } 2 \mid m \text{ and } 2 \nmid n \\ 1 & \text{if } 2 \mid m, \quad 2 \parallel n \\ -1 & \text{if } 2 \mid m, \quad 4 \mid n. \end{cases}$$

$$(7c) \quad B_{2^3} = \begin{cases} 0 & \text{if } 2 \nmid m, \text{ or } 4 \mid m, \text{ or } 2 \parallel m \text{ and } 4 \nmid n \\ 1 & \text{if } 2 \parallel m, \quad 4 \parallel n \\ -1 & \text{if } 2 \parallel m, \quad 8 \mid n. \end{cases}$$

$$q = 2^\lambda, \quad \lambda > 3. \quad (ii)$$

$$(7d) \quad B_{2^\lambda} = \begin{cases} 0 & \text{if } 2 \nmid m, \text{ or } 2 \parallel m, \\ & \text{or } 4 \mid m, 2 \nmid \lambda, 2^{\lambda-1} \nmid \Delta, \\ & \text{or } 4 \mid m, 2 \mid \lambda, 2^\lambda \nmid \Delta. \\ \frac{-C \cdot 2}{2^{(\lambda-1)/2}} & \text{if } 4 \mid m, 2 \nmid \lambda, 2^{\lambda-1} \mid \Delta; \\ \frac{-A \cdot 2}{2^{\lambda/2}} & \text{if } 4 \mid m, 2 \mid \lambda, 2^\lambda \mid \Delta. \end{cases}$$

where

$$A = \begin{cases} 1 & \text{if } \Delta/2^\lambda \equiv 0, 3 \pmod{4} \\ -1 & \text{if } \Delta/2^\lambda \equiv 1, 2 \pmod{4}. \end{cases}$$

$$C = \begin{cases} -1 & \text{if } \Delta/2^{\lambda-1} \equiv 3 \pmod{8} \\ 1 & \text{if } \Delta/2^{\lambda-1} \equiv 7 \pmod{8} \\ 0 & \text{all other cases.} \end{cases}$$

$$q = p^\lambda.$$

$$(7e) \quad B_{p^\lambda} = \begin{cases} 0 & \text{if } 2 \mid \lambda, p^{\lambda-1} \nmid \Delta, \\ & \text{or if } 2 \nmid \lambda \text{ and either } p^{\lambda-1} \nmid \Delta \text{ or } p^\lambda \mid \Delta. \\ \frac{-p^{(\lambda-2)/2}}{p^\lambda} & \text{if } 2 \mid \lambda, p^{\lambda-1} \parallel \Delta \\ \frac{p^{(\lambda-2)/2}(p-1)}{p^\lambda} & \text{if } 2 \mid \lambda, p^\lambda \mid \Delta \\ \left(\frac{-p^{-\lambda+1}\Delta}{p} \right) \frac{p^{(\lambda-1)/2}}{p^\lambda} & \text{if } 2 \nmid \lambda, p^{\lambda-1} \parallel \Delta. \end{cases}$$

Equations (7a) to (7d) give the following results:

If $2^\alpha \parallel \Delta$, $2^{-\alpha}\Delta = \Delta_1$, and $\alpha \geq 0$ is an integer, then

$$(8) \quad \chi_2(n, m) = \begin{cases} 0 & \text{if } m \not\equiv n \pmod{2} \\ 2 & \text{if } m \equiv n \equiv 1 \pmod{2} \\ 3 & \text{if } 2 \mid m, 2 \parallel n. \\ 1 - (-1)^{n/4} & \text{if } 2 \parallel m, 4 \mid n, \\ 3 \cdot 2^{(3-\alpha)/2} & \text{if } m \equiv n \equiv 0 \pmod{4}, \alpha \text{ is odd and } > 4 \\ E \cdot 2^{-\alpha/2} & \text{if } m \equiv n \equiv 0 \pmod{4}, \alpha \text{ is even and } \geq 4. \end{cases}$$

where

$$E = \begin{cases} 0 & \text{if } \Delta_1 \equiv 7 \pmod{8} \\ 4 & \text{if } \Delta_1 \equiv 3 \pmod{8} \\ 6 & \text{if } \Delta_1 \equiv 1, 5 \pmod{8}. \end{cases}$$

For each odd prime p , let β_p be the non-negative integer for which $p^{\beta_p} \parallel \Delta$ and let

$$\Delta_2 = \prod_{\substack{p \mid \Delta \\ \beta_p \equiv 1 \pmod{2}}} p^{\beta_p}, \quad \Delta_3 = \prod_{\substack{p \\ \beta_p \equiv 0 \pmod{2}}} p^{\beta_p}, \quad D = \prod_{p \mid \Delta_2} p.$$

Then Lomadze's results give:

$$(9) \quad \left\{ \begin{aligned} \varrho_4(n, m) &= \frac{4}{\pi} \Delta^{1/2} \chi_2 \prod_{p|D_2} \frac{1-p^{-(\beta_p+1)/2}}{1-p^{-1}} \\ &\prod_{p|D_2} \left[\frac{1-p^{-\beta_p/2}}{1-p^{-1}} \left\{ 1 - \left(\frac{-2^\alpha D}{p} \right) p^{-1} \right\} + p^{-\beta_p/2} \right] \sum_{\substack{q=1 \\ (q, 4D)=1}}^{\infty} \frac{(-2^\alpha D/q)}{p}. \end{aligned} \right.$$

That the infinite series in (9) is convergent and replaceable by a finite sum is shown in lemma 1 below.

Lemma 1. If D is a positive, odd, square-free integer and α a non-negative integer, then:

$$(10) \quad \sum_{\substack{q=1 \\ (q, 4D)=1}}^{\infty} \left(\frac{-2^\alpha D}{q} \right) \frac{1}{q} = \begin{cases} \frac{-\pi}{(8D)^{1/2}} \sum_{r=1}^{8D-1} \left(\frac{-8D}{r} \right) r & \text{if } 2 \nmid \alpha \\ \left[1 - \left(\frac{-D}{2} \right) \frac{1}{2} \right] \frac{-\pi}{D^{3/2}} \sum_{r=1}^{D-1} \left(\frac{-D}{r} \right) r & \text{if } 2 \mid \alpha \text{ and } D \equiv 3 \pmod{4} \\ \frac{-\pi}{(4D)^{3/2}} \sum_{r=1}^{4D-1} \left(\frac{-4D}{r} \right) r & \text{if } 2 \mid \alpha \text{ and } D \equiv 1 \pmod{4}. \end{cases}$$

Proof. If d is any integer such that $d \equiv 0$ or $1 \pmod{4}$ and not a square, we may define (cf. LANDAU [2, p. 145])

$$K(d) = \sum_{n=1}^{\infty} \left(\frac{d}{n} \right) \frac{1}{n}.$$

If d is further a fundamental discriminant number and $d < 0$, we have:

$$K(d) = \frac{-\pi}{|d|^{3/2}} \sum_{r=1}^{|d|-1} \left(\frac{d}{r} \right) r.$$

(cf. LANDAU [2, p. 172 and p. 177]).

Now, D being odd, $-8D$ is always a fundamental discriminant number; if $D \equiv 1 \pmod{4}$, $-4D$ is a fundamental discriminant number; and if $D \equiv 3 \pmod{4}$ then $-D$ is a fundamental discriminant number and:

$$K(-D) [(1 - (-D/2)^{1/2})] = K(-4D),$$

by Satz 214 of LANDAU [2, p. 172].

The proof of equation (10) follows.

From (8) and (9) we get an estimate for $\varrho_4(n, m)$.

$$|\kappa_2(n, m)| \leq 3,$$

$$\prod_{p|D_2} \frac{1-p^{-(\beta_p+1)/2}}{1-p^{-1}} < \prod_{p|D_2} \frac{1}{1-p^{-1}} \leq \max [1, N \log |D_2|],$$

where N is a positive numerical constant (cf. HARDY and WRIGHT [1, § 22.9]).

Similarly

$$\prod_{p|A_3} \left[\frac{1-p^{-\beta_p/2}}{1-p^{-1}} \left(1 - \left(\frac{-2^\alpha D}{p} \right) \frac{1}{p} \right) + p^{-\beta_p/2} \right] \leq \max [1, N^2 \log^2 |A_3|]$$

and

$$\left| \sum_{q=1}^{\infty} \left(\frac{-2^\alpha D}{q} \right) \frac{1}{q} \right| < \log |8D|$$

(cf. BATEMAN [1, p. 85]).

From the above we have ($\varepsilon > 0$)

$$(11) \quad \varrho_s(n, m) = O(\Delta^{1/s+\varepsilon}).$$

We now define $\theta(v|\tau)$ thus:

$$(12) \quad \theta(v|\tau) = \sum_n \sum_{\substack{m \\ m^2 \leq ns}} \varrho_s(n, m) w^n e^{2\pi i v m},$$

where in the above equation, as in the rest of this paper, we assume:

$$\begin{aligned} w &= e^{\pi i \tau} \\ \tau &= \alpha + i\beta \\ v &= v_1 + i v_2, \end{aligned}$$

with α, β, v_1 and v_2 real and $\beta > 0$.

We decompose $\theta(v|\tau)$ into the auxiliary functions $\psi(v|\tau)$ and $\vartheta_0(v|\tau)$ as follows:

$$(13) \quad \theta(v|\tau) = \vartheta_0(v|\tau) + \psi(v|\tau)$$

where

$$(14) \quad \vartheta_0(v|\tau) = \sum_{\substack{m \\ m \equiv 0(s)}} w^{m^2/s} e^{2\pi i v m}$$

$$(15) \quad \psi(v|\tau) = \sum_n \sum_{\substack{m \\ m^2 < ns}} \varrho_s(n, m) w^n e^{2\pi i v m}.$$

(13) follows readily from (12), (14) and (15) when we recall (1). $\vartheta_0(v|\tau)$ is clearly analytic and regular in v and τ ($\mathcal{J}(\tau) > 0$). Using (11) we see that the right hand side of (15) is majorized by

$$C_0(\varepsilon) \sum_{n=1}^{\infty} e^{-\pi n \beta} \cdot n^{1-\varepsilon} \cdot e^{2\pi |v_2| \sqrt{ns}} \quad (\varepsilon > 0, \beta > 0),$$

where $C_0(\varepsilon)$ is real, positive and a function of ε alone. Hence $\psi(v|\tau)$ and therefore $\theta(v|\tau)$ are analytic functions of v and τ , regular for $\mathcal{J}(\tau) > 0$.

It is convenient to replace the summation variable n by $k = ns - m^2$

in equation (15). Doing so and substituting for $\varrho_s\left(\frac{k+m^2}{s}, m\right)$ from (2) we obtain:

$$(16) \quad \left\{ \begin{aligned} \psi(v|\tau) &= \eta_s \sum_m w^{m^2/s} e^{2\pi i v m} \sum_{\substack{k=1 \\ k \equiv -m^2(s)}} k^{(s-3)/2} w^{k/s} \\ &\quad \sum_{q=1}^{\infty} \sum'_{a(q)} e\left(\frac{-ka}{qs}\right) \sum_{b(q)} \left(\frac{S(a, q, b)}{q}\right)^s e\left(\frac{-m^2 a - m b s}{qs}\right). \end{aligned} \right.$$

This may be re-written as:

$$(17) \quad \psi(v|\tau) = \sum_m w^{m^2/s} e^{2\pi i v m} \psi_m(\tau)$$

where

$$(18) \quad \psi_m(\tau) = \eta_s \sum_{\substack{k=1 \\ k \equiv -m^2(s)}}^{\infty} k^{(s-3)/2} w^{k/s} \sum_{q=1}^{\infty} \sum'_{a(q)} e\left(\frac{-ka}{qs}\right) \sum_{b(q)} \left(\frac{S(a, q, b)}{q}\right)^s e\left(\frac{-m^2 a - m b s}{qs}\right).$$

Now let

$$(19) \quad \sigma(a, q, m) = q^{(s-1)/2} \sum_{b(q)} \left(\frac{S(a, q, b)}{q}\right)^s e\left(\frac{-m^2 a - m b s}{qs}\right).$$

This definition of $\sigma(a, q, m)$ is formally identical with that of Bronkhorst who has developed its properties [1, pp. 11–13] and his work is easily seen to be valid when $s=4$.

We use these results beginning with the fact that $\sigma(a, q, m)$ has period s in m . That $\psi_m(\tau)$ also has period s in m follows when it is written in the form:

$$(20) \quad \psi_m(\tau) = \eta_s \sum_{\substack{k=1 \\ k \equiv -m^2(s)}}^{\infty} k^{(s-3)/2} w^{k/s} \sum_{q=1}^{\infty} q^{(1-s)/2} \sum'_{a(q)} e\left(\frac{-ka}{qs}\right) \sigma(a, q, m).$$

Hence $\psi(v|\tau)$ can be written as:

$$(21) \quad \left\{ \begin{aligned} \psi(v|\tau) &= \sum_{\mu(s)} \psi_{\mu}(\tau) \sum_{m \equiv \mu(s)} w^{m^2/s} e^{2\pi i v m} \\ &= \sum_{\mu(s)} \psi_{\mu}(\tau) \vartheta_{\mu}(v|\tau) \end{aligned} \right.$$

where (consistent with (14))

$$(22) \quad \vartheta_{\mu}(v|\tau) = \sum_{m \equiv \mu(s)} w^{m^2/s} e^{2\pi i v m}.$$

From the above we obtain easily

$$(23) \quad \theta(v|\tau) = \vartheta_0(v|\tau) + \sum_{\mu(s)} \psi_{\mu}(\tau) \cdot \vartheta_{\mu}(v|\tau).$$

To investigate the series in (18) we write it as: (cf. (4))

$$(24) \quad \psi_m(\tau) = \eta_s \sum_{\substack{k=1 \\ k \equiv -m^2(s)}}^{\infty} w^{k/s} k^{(s-3)/2} \sum_{q=1}^{\infty} B_q\left(\frac{k+m^2}{s}, m\right).$$

Since, when $s = 4$ and $p \nmid k$

$$B_p = \left(\frac{-k}{p}\right) \frac{1}{p}.$$

The inner sum in (24) is not absolutely convergent.

For $s > 5$ however the double series in (24) can be shown to be absolutely convergent and in this case one can reverse the order of summation in (24) obtaining:

$$(25) \quad \psi_m(\tau) = \eta_s \sum_{q=1}^{\infty} \sum'_{a(q)} q^{(1-s)/2} \cdot \sigma(a, q, m) \sum_{\substack{k=1 \\ k \equiv -m^2(s)}}^{\infty} e\left(\frac{-ka}{qs}\right) k^{(s-3)/2} w^{k/s} \quad (s > 5).$$

Using a formula of LIPSCHITZ [1] on the series in k , BRONKHORST [1, pp. 9–10] shows that:

$$(26) \quad \left\{ \begin{aligned} \psi_m(\tau) &= \sum_{q=1}^{\infty} \sum'_{a(q)} q^{(1-s)/2} \sigma(a, q, m) s^{-1/2} e\left(\frac{s-1}{8}\right) 2^{-(s-1)/2} \\ &\quad \sum_r \frac{e(-m^2 r/s)}{(\tau/2 - a/q - r)^{(s-1)/2}}, \end{aligned} \right.$$

where

$$0 < \arg \sqrt{\tau/2 - a/q - r} < \pi/2.$$

While the above transformations have not been justified for $s=4$, they motivate our definition of the function $\psi_{m,z}(\tau)$ below.

Let $z = \sigma + it$ and put:

$$(27) \quad \left\{ \begin{aligned} \psi_{m,z}(\tau) &= \frac{s^{-1/2} e\left(\frac{s-1}{8}\right)}{2^{(s-1)/2}} \sum_{q=1}^{\infty} \frac{2^{\delta(q)z}}{q^{(s-1)/2+z}} \sum'_{a(q)} \sigma(a, q, m) \\ &\quad \sum_r \frac{e(-m^2 r/s)}{\left(\frac{\tau}{2} - \frac{a}{q} - r\right)^{(s-1)/2}} 2^z \left| \frac{\tau}{2} - \frac{a}{q} - r \right|^z \quad (s=4, \sigma > 1/2). \end{aligned} \right.$$

It will now be shown that $\psi_{m,z}(\tau)$ is analytic in z for $\Re(z) > 1/2$ (v and τ fixed).

Making the change of variable $a' = a + rq$ and using the periodicity of $S(a, q, b)$ in a it is easily shown (cf. BRONKHORST [1, p. 10]) that:¹⁾

$$(28) \quad \psi_{m,z}(\tau) = s^{-1/2} e\left(\frac{s-1}{8}\right) \sum_{q=1}^{\infty} \sum_{\substack{a \\ (a,q)=1}} \frac{2^{\delta(q)z} \sigma(a, q, m)}{(q\tau - 2a)^{(s-1)/2} |q\tau - 2a|^z}.$$

We need estimates for $\sigma(a, q, m)$. From Bronkhorst's results (pp. 11–13) we have:

$$(29) \quad |\sigma(a, q, m)| = \begin{cases} q^{-1/2} |S(-as, q, -2am)|, & \text{if } q \equiv 1 \pmod{2} \\ (q/2)^{-1/2} 2^{(s-1)/2} |S(-2as, q/2, -2am)|, & \text{if } q \equiv 2 \pmod{4} \\ (q)^{-1/2} |S(-as, q, -2am)| 2^{(s-2)/2}, & \text{if } q \equiv 0 \pmod{4}. \end{cases}$$

¹⁾ The inner sum in (28) is absolutely convergent for $\Re(z)$ sufficiently large.

By well-known bounds for $|S(a, q, b)|$ it now follows that:

$$(30) \quad |\sigma(a, q, m)| \leq 2^{(s-1)/2}.$$

Thus the series defining $\psi_{m,z}(\tau)$ in (28) is majorized by:

$$(31) \quad 2^{(s-1)/2} s^{-1/2} \sum_{\substack{q=1 \\ (a,q)=1}}^{\infty} \sum_a \frac{2^{\delta(q)\sigma}}{|q\tau-2a|^{(s-1)/2+\sigma}}.$$

which converges for $\frac{s-1}{2} + \sigma > 2$ (cf. L. R. FORD [1, pp. 151-52]). Hence when $s=4$, the series converges uniformly for $\sigma > 1/2$, so that $\psi_{m,z}(\tau)$ is a regular, analytic function of z in the region: $\Re(z) > 1/2$.

Since $\psi_{m,z}(\tau)$, like $\psi_m(\tau)$, has periodicity s in m , we define:

$$(32) \quad \psi_z(v|\tau) = \sum_{\mu \bmod s} \psi_{\mu,z}(\tau) \cdot \partial_{\mu}^{\mathbf{v}}(v|\tau) \quad (\Re(z) > 1/2),$$

$$(33) \quad \theta_z(v|\tau) = \partial_0(v|\tau) + \psi_z(v|\tau), \quad (\Re(z) > 1/2).$$

We proceed to show that $\psi_{m,z}(\tau)$ (and hence also $\psi_z(v|\tau)$ and $\theta_z(v|\tau)$) can be continued analytically into the region $\Re(z) > -1/2$ and that:

$$\psi_{m,z}(\tau)|_{z=0} = \psi_{m,0}(\tau) = \psi_m(\tau).$$

In equation (27) we consider the inner sum and define:

$$(34) \quad P' = \sum_r \frac{e(-m^2 r/s)}{(\tau/2 - a/q - r)^{(s-1)/2} |\tau/2 - a/q - r|^z}.$$

Thus

$$(35) \quad \psi_{m,z}(\tau) = \frac{s^{-1/2} e\left(\frac{s-1}{8}\right)}{2^{(s-1)/2+z}} \sum_{q=1}^{\infty} \sum_{a(q)}' \frac{\sigma(a, q, m) 2^{\delta(q)z}}{q^{(s-1)/2+z}} P' \quad (\Re(z) > 1/2).$$

Now, (34) is unaltered when $-r$ is replaced by r . We do this and apply to (34) the Poisson summation formula (L. J. MORDELL [2]) to obtain:

$$P' = \sum_n \int_{-\infty}^{\infty} \frac{e\left(\frac{m^2 x}{s}\right) e(-nx) dx}{(\tau/2 - a/q + x)^{(s-1)/2} |\tau/2 - a/q + x|^z}.$$

On writing:

$$k = ns - m^2, \quad u = x - a/q,$$

We have:

$$(36) \quad \begin{cases} P' = \sum_{k=-m^2(s)}^{\infty} \int_{-\infty}^{\infty} \frac{e(-ku/s) \cdot e(-ka/qs) du}{(\tau/2 + u)^{(s-1)/2} |\tau/2 + u|^z} \\ = \sum_{k=-m^2(s)}^{\infty} e(-ka/qs) \cdot I(k, \tau, z), \end{cases}$$

where (36) defines $I(k, \tau, z)$ in an obvious way.

In what follows S will denote an arbitrary but fixed compact subset of the region $\mathcal{R}(z) > -1/2$, containing the set $\{z: -1/4 < \sigma < 1, |t| < 1\}$. The constants $C_1, C_2, \dots C_i \dots$ are assumed to be positive and depend at most on S and τ both of which are assumed to be fixed for the rest of this section, $\mathcal{I}(\tau)$ being positive.

Lemma 2. The function $I(k, \tau, z)$ has the following properties:

- a) For each k and τ ($\mathcal{I}(\tau) > 0$), $I(k, \tau, z)$ is an analytic function of z , regular for $\mathcal{R}(z) > -1/2$.
- b) In the set S ,

$$|I(k, \tau, z)| < C_1 e^{-C_2 |k|}.$$

Lemma 3.

$$(37) \quad I(k, \tau, 0) = \begin{cases} 0 & \text{if } k \leq 0, \\ \frac{e\left(\frac{1-s}{8}\right) (2\pi)^{(s-1)/2} (k/s)^{(s-3)/2} e^{\pi i \tau k}}{\Gamma\left(\frac{s-1}{2}\right)} & \text{if } k > 0. \end{cases}$$

Lemmas 2 and 3 appear in almost the present form in the thesis of H. STREEFKERK [1, p. 31]. The method of proof was indicated by HECKE [1, pp. (224–226)]. See also part I of same paper (p. 111). A detailed proof can be found in SESHU¹⁾, [1, p. 24].

From (35) and (36) we get:

$$(38) \quad \left\{ \begin{aligned} \psi_{m,z}(\tau) &= \frac{s^{-1/2} e\left(\frac{s-1}{8}\right)}{2^{(s-1)/2+z}} \sum_{q=1}^{\infty} \sum_{a(q)} \frac{\sigma(a, q, m) 2^{\delta(q)z}}{q^{(s-1)/2+z}} \\ &\quad \sum_{k=-m^2(s)} I(k, \tau, z) e\left(\frac{-ka}{qs}\right), \quad (\sigma > 1/2). \end{aligned} \right.$$

From the estimates for $I(k, \tau, z)$ in lemma 2b), the double series in (38) is absolutely convergent for $\mathcal{R}(z) > 1/2$. On interchanging summations and using (19) and (4) we get (taking, as usual, $\frac{k+m^2}{s} = n$) that:

$$(39) \quad \psi_{m,z}(\tau) = \frac{s^{-1/2} e\left(\frac{s-1}{8}\right)}{2^{(s-1)/2+z}} \sum_{k=-m^2(s)} I(k, \tau, z) \sum_{q=1}^{\infty} \frac{B_q(n, m) 2^{\delta(q)z}}{q^z}.$$

Define

$$(40) \quad G_z(n, m) = \sum_{q=1}^{\infty} \frac{B_q(n, m) 2^{\delta(q)z}}{q^z} \quad (\mathcal{R}(z) > 1/2).$$

From (7a) through (7e), it follows that:

$$|B_q(n, m)| < 4q^{-1/2}.$$

where

$$q = p^\lambda \text{ or } q = 2^\lambda, \quad (\lambda, a \text{ positive integer}).$$

¹⁾ See also BATEMAN [1].

Consequently one may use the multiplicativity of $B_q(n, m)$ to write:

$$(41) \quad G_z(n, m) = \chi_{2,z}(n, m) \cdot F_z(n, m) \quad (\mathcal{R}(z) > 1/2),$$

where

$$(42) \quad \chi_{2,z}(n, m) = 1 + 2^z \sum_{\lambda=1}^{\infty} B_{2^\lambda} 2^{-\lambda z}.$$

We now show that $G_z(n, m)$ is an analytic function, regular for $\mathcal{R}(z) > -1/2$ and under a bound, depending at most on k , in the set S , where, as before, $k = ns - m^2$.

Lemma 3. $F_z(n, m)$ is an analytic function of z which is:

- a) regular for $\mathcal{R}(z) > -1/2$ when $-k$ is not a square,
- b) regular in the half-plane $\mathcal{R}(z) > -1/2$ except at the point $z=0$, when $k=0$ or when $k < 0$ and $-k$ is a square. In these cases $F_z(n, m)$ has a simple pole at $z=0$.

Proof. For convenience we define:

$$(43) \quad \chi_{p,z}(n, m) = 1 + \sum_{\lambda=1}^{\infty} B_{p^\lambda}(n, m) p^{-\lambda z} \quad (\sigma > 1/2).$$

By (7e) $\chi_{p,z}(n, m)$ depends only on k and not on n and m separately¹⁾.

Inspection of (7e) shows that when $k \neq 0$, the right hand side of (43) has at most $\beta + 2$ non-zero terms if $p^\beta \parallel k$ (β being a non-negative integer). Thus $\prod_{p|k} \chi_{p,z}(k)$ is an entire function of z when $k \neq 0$.

Since

$$|B_{p^\lambda}(k)| < p^{-\lambda/2},$$

an easy calculation shows that:

$$(44) \quad |\chi_{p,z}(k)| < C_3 p^\beta \quad (\mathcal{R}(z) > -1/2).$$

When $k \neq 0$, (7e) gives:

$$(45) \quad \prod_{p \nmid k} \chi_{p,z}(k) = \prod_{p \nmid k} \left(1 + \left(\frac{-k}{p} \right) p^{-1-z} \right) \quad (\mathcal{R}(z) > 0).$$

By the multiplicativity of $B_{q,z}(n, m)$ one gets:

$$(46) \quad \left\{ \begin{aligned} F_z(n, m) &= \sum_{\substack{q=1 \\ q \equiv 1(2)}}^{\infty} B_q(n, m) q^{-z} = \prod_{p|k} \chi_{p,z} \cdot \prod_{p \nmid k} \chi_{p,z} \\ &= \prod_{p|k} \chi_{p,z} \cdot \prod_{p \nmid k} \left(1 + \left(\frac{-k}{p} \right) p^{-1-z} \right) \\ &= \prod_{p|k} \chi_{p,z} \cdot \sum_{q=1}^{\infty} \left(\frac{-4k}{q} \right) q^{-1-z} \mu^2(q) \quad (\mathcal{R}(z) > 0), \end{aligned} \right.$$

¹⁾ In the rest of the section, we write $B_{p^\lambda}(k)$ ($\chi_{p,z}(k)$) in place of $B_{p^\lambda}(n, m)$ ($\chi_{p,z}(n, m)$), whenever convenient.

where μ is the Möbius function. Now $\left(\frac{-4k}{q}\right)$ is a residue class character modulo $|4k|$ and is the principal character only if $-4k$ is a square, (LANDAU [2, satz. 99]). We denote this character (defined on the positive integers) by $\chi(q)$. STREEFKERK [1, hulpt. 24] has shown that:

$$(47) \quad \left\{ \begin{aligned} \sum_{q=1}^{\infty} \chi(q) \mu^2(q) q^{-1-z} &= L(1+z, \chi) \cdot \prod_p \left(1 - \frac{\chi(p^2)}{p^{2+2z}}\right) \\ &= L(1+z, \chi) \sum_{d=1}^{\infty} \frac{\mu(d) \chi(d^2)}{d^{2+2z}}. \end{aligned} \right.$$

When χ is not a principal character (i.e. $-k$ is not a square) $L(1+z, \chi)$ is regular for $\Re(z) > -1$;

$$\sum_{d=1}^{\infty} \frac{\mu(d) \chi(d^2)}{d^{2+2z}}$$

defines a regular analytic function for $\Re(z) > -1/2$. From $Eq.$'s (46), (47) and the principle of analytic continuation, $F_z(n, m)$ is regular for $\Re(z) > -1/2$.

When $-k$ is a square, ($k \neq 0$), χ is a principal character and $L(1+z, \chi)$ can be decomposed as:

$$(48) \quad L(1+z, \chi) = \prod_{p|k} (1 - p^{-1-z}) (1 - 2^{-1-z}) \zeta(1+z).$$

where ζ denotes the Riemann-Zeta function. (See LANDAU [1, p. 423]). Given the well-known properties of $\zeta(1+z)$, the equations (46) to (48) give the analytic continuation of $F_z(n, m)$ into the region $\Re(z) > -1/2$ and show that it is regular there except for (at most) a simple pole at $z=0$. That it does have a pole follows if the other factors of $F_z(n, m)$ have no zeros there. The only possibility is $\prod_{p|k} \chi_{p,z}(k)$: but when $-k$ is a square, $k \neq 0$, (7e) gives, if $p^{2b} \parallel k$, (b a positive integer).

$$\begin{aligned} \chi_{p,z}(k) \Big|_{z=0} &= \chi_p(k) = \left(1 + \frac{p-1}{p^2} + \frac{p-1}{p^3} + \dots + \frac{p-1}{p^{b+1}} + \frac{1}{p^{b+1}}\right) \\ &= 1 + 1/p. \end{aligned}$$

We have now to compute $F_z(n, m)$ when $k=0$. By (7e)

$$(49) \quad \left\{ \begin{aligned} \chi_{p,z}(0) &= 1 + \frac{p-1}{p^{2+2z}} + \frac{p-1}{p^{3+2z}} + \dots \\ &= \frac{1-p^{-2-2z}}{1-p^{-1-2z}} \quad (\sigma > -1/2). \end{aligned} \right.$$

If $\sigma > 0$, we have $F_z(n, m)$ defined by an absolutely convergent product:

$$(50) \quad \left\{ \begin{aligned} F_z(n, m) &= \prod_p \chi_{p,z}(0) = \prod_p \frac{1-p^{-2-2z}}{1-p^{-1-2z}} \\ &= \frac{1-2^{-1-2z}}{1-2^{-2-2z}} \frac{\zeta(1+2z)}{\zeta(2+2z)}. \end{aligned} \right.$$

The last member of (50) is regular for $\Re(z) > -1/2$ except for a simple pole at $z=0$ and this completes the proof of the lemma.

Lemma 5:

- a) $\chi_{2,z}(n, m)$ is an analytic function of z , regular at least in the region $\Re(z) > -1/2$ for all n, m .
- b) For those values of n and m for which $k=0$ or $-k$ is a square, $\chi_{2,z}(n, m)$ has a zero at $z=0$.

Proof. First consider $k \neq 0$. Inspection of (7a) to (7d) reveals that the series

$$\sum_{\lambda=1}^{\infty} B_{2\lambda}(n, m) 2^{-\lambda z}$$

has at most $\alpha+1$ terms, when $2^\alpha \parallel k$ (α being a non-negative integer). Each term is an entire function of z , thus also $\chi_{2,z}(n, m)$.

Since also

$$|B_{2\lambda}(n, m)| < 4 \cdot 2^{-\lambda/2},$$

it follows easily that

$$(51) \quad |\chi_{2,z}(n, m)| < C_5 2^\alpha, \quad (\sigma \geq -1/2, k \neq 0).$$

When $k=0$, $\chi_{2,z}(n, m)$ is not necessarily a finite sum of terms. It depends on the residue classes mod 4 or mod 8 to which n and m belong. The condition:

$$m^2 = 4n$$

restricts the possible n, m combinations to the following: (1) $2 \parallel m$, thus $2 \nmid n$ or (2) $4 \mid m$ and $4 \mid n$. If λ represents a positive integer, then from (7a) to (7d) we see that $B_{2\lambda}(n, m)$ is zero for all allowable pairs n, m except as listed below in (52):

$$(52) \quad B_{2\lambda}(n, m) = \begin{cases} (-1)^n & \text{if } \lambda = 1 \\ -2/2^{\lambda/2} & \text{if } 2 \mid \lambda, \lambda \geq 2 \\ & \text{and } 4 \mid m, 4 \mid n. \end{cases}$$

The corresponding expressions for $\chi_{2,z}(n, m)$ emerge as:

$$(53) \quad \chi_{2,z}(n, m) = \begin{cases} 0 & \text{if } 2 \parallel m, 2 \nmid n. \\ \frac{(2^z-1)(2^{1+z}+1)}{2^{2z}(1-2^{-1-2z})} & \text{if } 4 \mid m, 4 \mid n, \quad (\sigma \geq -1/2). \end{cases}$$

This completes the proof of a) and, in fact, part of b): for (53) shows that

$$\chi_{2,0}(n, m) = 0.$$

To continue with b): when $-k$ is a square ($k < 0$), the possible combinations of n, m are again restricted. $B_{2\lambda}(n, m)$ can be evaluated from (7a) to (7d) for all permissible combinations (see Eq. (54) for the

possibilities) and $\chi_{2,z}(n, m)$ calculated thereby. The expressions for $\chi_{2,z}(n, m)$ after simplification, are:

$$(54) \quad \chi_{2,z}(n, m) = \begin{cases} 0 & \text{if } 2 \nmid m, 2 \mid n \text{ or if } \\ & 2 \parallel m, 2 \nmid n \text{ or if } 4 \mid m, 2 \nmid n, & (a) \\ 2^{-2z}(2^{1+2z} + 1)(2^z - 1) & \text{if } 2 \parallel m, 8 \mid n, & (b) \\ \frac{(2^{a+2az}-1)(2^{1+z}+1)(2^z-1)}{(2^{1+2z}-1)2^{a-1+2az}} & \text{if } 4 \mid m, 4 \mid n & (c), \end{cases}$$

where, in (c), a is an integer such that $2^{2a} \parallel k$; clearly $a \geq 2$.

As before the factor $2^z - 1$ ensures that:

$$\chi_{2,0}(n, m) = 0,$$

when $k \neq 0$, and $-k$ is a square.

Lemma 6. a) The function $G_z(n, m)$ has an analytic continuation into the half-plane $\Re(z) > -1/2$ which is regular there.

b) When $-k$ is not a square,

$$(55) \quad G_z(n, m)|_{z=0} = \sum_{q=1}^{\infty} B_q(n, m).$$

Proof. Part a) is an immediate consequence of equation (41) and lemmas 4 and 5.

To prove b) we observe that when χ is non-principal (mod $|4k|$)

$$L(1, \chi) = \sum_{j=1}^{\infty} \frac{\chi(j)}{j} = \sum_{j=1}^{\infty} \left(\frac{-4k}{j} \right) j^{-1},$$

where the series on the right is convergent (LANDAU [2, satz. 141]). Thus the equation (obtained from (47) by letting $z=0$):

$$\sum_{q=1}^{\infty} \chi(q) \mu^2(q)/q = \sum_{d=1}^{\infty} \frac{\mu(d) \chi(d^2)}{d^2} \sum_{j=1}^{\infty} \frac{\chi(j)}{j}$$

has more than formal significance: for the left member of the equation is the Dirichlet product of the absolutely convergent and the convergent series occurring on the right hand side.

Thus from (46) and (41):

$$G_z(n, m)|_{z=0} = \chi_2(n, m) \prod_{p \mid k} \chi_p(k) \sum_{\substack{q=1 \\ (q, 4k)=1}}^{\infty} B_q(n, m) = \sum_{q=1}^{\infty} B_q(n, m).$$

Lemma 7. In the set S ,

$$(56) \quad |G_z(n, m)| < C_6 k^2.$$

Proof. When $k=0$, $\chi_{2,z}(n, m)$ and $F_z(n, m)$ are given by (53) and (50) respectively. The product is regular in S and independent of k, n or m : thus

$$(57) \quad |G_z(n, m)| < C_7 \quad (k=0).$$

When $-k$ is not a square, χ is non-principal and thus ¹⁾:

$$(58) \quad |L(1+z, \chi)| < C_8 |k|.$$

We next observe that

$$\sum_{d=1}^{\infty} \frac{\mu(d) \chi(d^2)}{d^{2+2z}}$$

defines a function independent of k and regular, thus bounded, in S . From equations (41), (46) and (47) and inequalities (44) and (51) (valid when $k \neq 0$) we have:

$$(59) \quad |G_z(n, m)| < C_9 k^2, \quad (-k \text{ non-square}).$$

When $k < 0$ and $-k$ is a square $L(1+z, \chi)$ is given by (48) and $\chi_{2,z}(n, m)$ by (54). Thus the product $\chi_{2,z}(n, m) \cdot \zeta(1+z)$ is bounded in S . Since further

$$(60) \quad \prod_{p|k} |(1-p^{-1-z})| < \prod_{p|k} p, \quad (\sigma \geq -1/2)$$

and

$$|(2^{a+2az}-1) 2^{1-a-2az}| < 4 \quad (\sigma \geq -1/2),$$

it easily follows that

$$(61) \quad |G_z(n, m)| < C_{10} k^2, \quad (k < 0, -k \text{ a square})$$

This completes the proof of lemma 7.

Theorem 1. The functions $\psi_{m,z}(\tau)$, $\psi_z(v|\tau)$ and $\theta_z(v|\tau)$ can be continued analytically into the half-plane $\Re(z) > -1/2$. All three functions are regular in this region; and at $z=0$, they are respectively identical with $\psi_m(\tau)$, $\psi(v|\tau)$ and $\theta(v|\tau)$ as defined in (24), (17) and (12).

Proof. Recall from (39) and (40) that:

$$(62) \quad \psi_{m,z}(\tau) = \frac{s^{-1/2} e\left(\frac{s-1}{8}\right)}{2^{(s-1)/2+z}} \sum_{k=-m^2(s)} I(k, \tau, z) G_z(n, m) \quad (\Re(z) > 1/2).$$

By lemmas 2 and 6, the terms of the series in (62) are functions regular in z for $\Re(z) > -1/2$. From lemmas 2 and 7 it follows that the series:

$$C_{11} + \sum_{k \neq 0} C_{12} k^2 \cdot e^{-|k| C_2}$$

majorizes the series in (62) in set S . Thus the latter series defines an analytic function of z regular in S . Since S is an arbitrary set in the region $\Re(z) > -1/2$, containing the neighbourhood

$$\{z : -1/2 < \sigma < 1, |t| < 1\},$$

$\psi_{m,z}(\tau)$ has an analytic extension regular for $\Re(z) > -1/2$.

¹⁾ See LANDAU [2, satz 235].

The series in (62) is uniformly convergent in S ; thus on using lemmas 3 and 6 we have (by (24)):

$$(63) \quad \left\{ \begin{aligned} \psi_{m,0}(\tau) &= \frac{\pi^{(s-1)/2}}{s^{(s-2)/2} \cdot \Gamma\left(\frac{s-1}{2}\right)} \sum_{\substack{k > 0 \\ k \equiv -m^2(s)}} e^{\pi i \tau k} \cdot k^{(s-3)/2} \sum_{q=1}^{\infty} B_q((k+m^2)/s, m) \\ &= \psi_m(\tau). \end{aligned} \right.$$

From (32) and (33) we get at once:

$$(64) \quad \psi_0(v|\tau) = \psi_z(v|\tau)|_{z=0} = \psi(v|\tau)$$

$$(65) \quad \theta_0(v|\tau) = \theta_z(v|\tau)|_{z=0} = \theta(v|\tau).$$

(To be continued)